


## Sundew ' Drosera trinervia' (CarniDr91a)

## Drosera trinervia

€ 8.29

Drosera trinervia is a widely spread South African carnivorous plant. It earns its name to the three veins that run trough the underside of each trapping leaf. The sundew is dormant in the summer and re-emerges in the winter.





## Product info

Drosera trinervia is a South African sundew that has a large area of distribution. On the underside of each of the trapping leaves three veins run. This characteristic has earned Drosera trinervia its name. This carnivorous plant is dormant in the summer and will die of down to the roots in the warmer period of the year. In the winter the sundew will re-emerge above the surface.

Height: 4 cm	<b>#</b> Ī	Width: 5 cm	
Adult height: 5-10 cm	<b>♣</b> <u>Ī</u>	Adult width: 10 cm	
Flower period: Jan-July	<b>⊕</b>	Product: plant	÷
Plant type: carnivorous		Growth habit: normal	*>
Properties: special	34	Growth rate: normal	<u></u>
Araflora info: introduction	wife of toth	Natural habitat: South Africa	800
Altitude range: 0-500 m	<u>₹</u>		

## Plantcare

after 2 years

Drosera trinervia can be cultivated in a mixture of peat, sand and perlite and is best kept quite moist during its active growing period in winter. When it gets warmer in the summer Drosera trinervia will go into a dormant state. When this happens the soil can be left to dry slightly, leaving it moderately moist.

Care: normal	<b>\$</b>	Place: indoor	圕
Place: artificial lighting	•	Light: full light	
Artificial light height: 0-20cm		Fertilize: none	NPK ,
Maintenance: prune dead stalks	À	Watering: high	
Watering tips: standing in water		Watering tips: no-water spraying	X
Air humidity at day: 60-100%		Air humidity at night: 80-100%	<b>&amp;</b>
Air movement: normal	<b>*</b> *	Temperature 15°C	
Day temperature 15-25°C	<b>₩</b>	Night temperature 10-20°C	<b>€</b> MAX
Temperature max: 30°C	<b>I</b>	Soil type: peat	
Repotting: after 2 years			